

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

AD-A235 743

2

REPORT DO




1a. REPORT SECURITY CLASSIFICATION UNCL		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2a. SECURITY CLASSIFICATION AUTHORITY DTIC		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE MAY 23 1991			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMRI 90-143			
6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research Institute	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Naval Medical Command	
6c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20889-5055		7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, DC 20372-5120	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Medical Research & Development Command	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20889-5044		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 61102A	PROJECT NO. BM161102.BS13
		TASK NO. AK111	WORK UNIT ACCESSION NO. DA313955
11. TITLE (Include Security Classification) A malaria sporozoite surface antigen distinct from the circumsporozoite protein			
12. PERSONAL AUTHOR(S) Hedstrom RC, Campbell JR, Leef ML, Charoenvit Y, Carter M, Sedegah M, Beaudoin RL, Hoffman SL			
13a. TYPE OF REPORT journal article	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) 1990	15. PAGE COUNT 6
16. SUPPLEMENTARY NOTATION Reprinted from: Bulletin of the World Health Organization 1990 Vol.68 (Supplement) pages 152-157			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		malaria vaccine; sporozoite mRNA; sporozoite surface antigen; polymerase chain reaction; monoclonal antibody; keyhole limpet hemocyanin; Plasmodium yoelii; circumsporozoite protein; antigen genes	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<div style="text-align: center;"> </div> <div style="float: right;"> <p>Accession for</p> <p>DTIC GRA&amp;I</p> <p>DTIC TAB</p> <p>Unannounced</p> <p>Justification</p> <p>By</p> <p>Distribution/</p> <p>Availability Codes</p> <p>Dist Avail and/or Special</p> <p>A-1 20</p> </div>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Phyllis Blum, Librarian		22b. TELEPHONE (Include Area Code) (301) 295-2188	22c. OFFICE SYMBOL MRL/NMRI

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.  
All other editions are obsolete.SECURITY CLASSIFICATION OF THIS PAGE  
UNCLASSIFIED

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## A malaria sporozoite surface antigen distinct from the circumsporozoite protein

R.C. Hedstrom,<sup>1,2</sup> J.R. Campbell,<sup>1,3</sup> M.L. Leef,<sup>1</sup> Y. Charoenvit,<sup>1</sup> M. Carter,<sup>4</sup> M. Sedegah,<sup>1</sup> R.L. Beaudoin,<sup>1</sup> & S.L. Hoffman<sup>1</sup>

*Monoclonal antibody NYS4 recognizes a single 140 kDa antigen on the surface of Plasmodium yoelii sporozoites, an antigen which is distinct from the extensively characterized circumsporozoite (CS) protein. To more thoroughly characterize this additional surface component, a genomic expression library was screened with NYS4 and an immunoreactive clone (M4) was obtained which expressed part of the antigen gene. The deduced amino acid sequence of the M4 peptide included two unique repetitive sequences of amino acids and a conserved sequence motif which is found in several proteins including the CS protein (region II). The cloned DNA hybridized to a PCR (polymerase chain reaction) amplified sporozoite mRNA demonstrating the sporozoite-stage expression of this gene. A synthetic peptide of one of the repeats, (Asn-Pro-Asn-Glu-Pro-Ser), was recognized by NYS4 and mice immunized with (Asn-Pro-Asn-Glu-Pro-Ser)<sub>3</sub> conjugated to KLH (keyhole limpet haemocyanin) produced high levels of antibodies that reacted with the surface of sporozoites and specifically to the 140 kDa antigen. Thus, at least two different proteins are on the surface of the P. yoelii sporozoite indicating that the immunoreactive exterior of the infective stage of malaria parasites is more antigenically complex than previously thought.*

### Introduction

The sporozoite of *Plasmodium* is uniformly covered by a proteinaceous membrane coat which is thought to be composed entirely of the circumsporozoite (CS) protein (1, 2). Being the first parasite molecule encountered by the host, the CS protein has been intensively studied and evaluated as a vaccine against malaria. Indeed, experimental immunization with radiation-attenuated sporozoites induces solid protective immunity (3-5) coincident with the stimulation of both cellular and humoral immunological responses to the CS protein (6, 7). An emerging view is that an essential feature of the protective response induced by sporozoites is cellular immunity (4-9). If this is correct, then a multicomponent vaccine containing additional pre-erythrocytic stage antigens may be required to induce sterilizing immunity in all individuals, since those regions of the CS protein that are the immunodominant determinants of T-cell responses are also the most variable in amino acid sequence from parasite to parasite (10-12).

The search for additional antigens by Charoenvit et al. led to the production of several monoclonal

antibodies that recognize sporozoite determinants which are distinct from the CS protein (13). One of these antibodies, designated NYS4 (Navy Yoelii Sporozoite 4), recognizes a sporozoite-specific 140 kDa antigen. This antigen is expressed on the surface of live sporozoites and is secreted along with the CS protein *in vitro* (unpublished results). Wortman et al. used this antibody to isolate an antigen-expressing clone from a *P. yoelii* genomic expression library (14). Here we describe the cloning and characterization of a portion of the gene encoding the 140 kDa antigen and identify a hexapeptide of this protein that is a B-cell epitope on sporozoites.

### Materials and methods

**Parasites and DNA isolation.** *P. yoelii* 17 X (NL) parasites were obtained by blood passage in Balb C mice. DNA isolation from parasite infected blood was performed, as described (14).

**Genomic expression library and immunoselection.** A *P. yoelii* genomic expression library was constructed using 0.5-2.0 kb fragments generated by partial DNase I digestion and commercial (Promega, Madison, WI) lambda gt11 arms and packaging extracts (15). The library was screened for antigen expressing clones with a 1:20 dilution of NYS4 (13) hybridoma culture supernatant, as described previously (16-18). NYS4 immunoreactive plaques were detected with a commercially prepared antibody detection kit

<sup>1</sup> Malaria Program, Infectious Diseases Department, Naval Medical Research Institute, Bethesda, MD 20814-5055, USA. Correspondence should be sent to Dr S.L. Hoffman at this address.

<sup>2</sup> U.S. Naval Medical Research Unit 3, Cairo, Egypt.

<sup>3</sup> Naval Research Laboratory, Washington, DC, USA.

<sup>4</sup> Department of Immunology, Walter Reed Army Institute of Research, Washington, DC, USA.

91-00295



Fig. 1 (a). Nucleotide sequence and derived amino acid sequence of a portion of the *P. yoelii* genomic DNA insert in lambda gMSY-4 (M4). The sequences of oligonucleotide primers (+) and (-) used for PCR amplification are indicated.

CCT TTT CTT ACT AAA GTT TGT CAG GAA GTA GAA AGA ATT	(+)----->	60
P P L T K V C Q E V E R I A H C G K W E		
GAA TGG AGT GAA TGT TCT ACT ACT TGT GAT GAA GGA AGA AAA ATT AGA AGA AGA CAA ATA		120
E W S E C S T T C D E G R A K I R R R Q I		
TTA CAT CCT GGA TGT GTT AGT GAG ATG ACT ACT CCA TGT AAG GTT CGT GAT TGC CCA CAA		180
L H P G C V S E M T T P C K V R D C P Q		
ATA CCA ATA CCT CCT GTC ATC CCT AAT AAA ATT CCA GAA AAG CCA TCA AAC CCA GAA GAA		240
I P I P P V I P N K I P E K P S N P E E		
CCA GTA AAT CCA AAC GAT CCA AAC GAT	-----3-MER REPEATS----->	300
P V N P N D P N D P N N P N N P N N P N N P N N P N N		
AAC CCA AAC AAC CCA AAT AAC CCA AAC AAC CCA AAC AAC CCA AAC AAC CCA AAC AAC CCA		360
N P N N P N N P N N P N N P N N P N N P N N P N N P N N P		
AAC AAT CCA AAT AAC CCA AAT AAC CCA AAC AAC CCA AAT AAC CCA AAT AAC CCA AAC AAC		420
N N P N N P N N P N N P N N P N N P N N P N N P N N P N N P		
CCA AAT AAC CCA AAC AAC CCA AAT AAC CCA AAT AAC CCA AAT AAC CCA AAT AAC CCA AAC	-----3-MER REPEATS----->	480
P N N P N N P N N P N N P N N P N N P N N P N N P N N P N N		
GAT CCA TCA AAC CCA AAC AAC CAC CCA AAA AGG CGA AAC CCA AAA AGG CGA AAC CCA AAC		540
D P S N P N N H P K R R P K R R N P N		
AAG CCA AAA CCA AAC AAG CCA AAC CCA AAC AAG CCA	-----6-MER REPEAT REGION----->	600
K P K P N K P N P N K P N P N P N E P S N P		
AAC AAG CCA AAC CCA AAC GAA CCA TCA AAC CCA AAC AAG CCA AAC CCA AAC GAA CCA TCA		660
N K P N P N E P S N P N K P N P N P N E P S		
AAC CCA AAC AAG CCA AAC CCA AAT GAG CCA TCA AAC CCA AAC AAG CCA AAC CCA AAT GAG		720
N P N K P N P N E P S N P N K P N P N P N E		
CCA TTA AAC CCA AAC GAG CCA TCA AAT CCA AAC GAG CCA TCA AAC CCA AAT GCG CCA TCA		780
P L N P N E P S N P N E P S N P N A P S		
AAC CCA AAC GAA CCA TCA AAC CCA AAT GAA CCA TCA AAC CCA AAT GAG CCA TCA AAC CCA		840
N P N E P S N P N E P S N P N E P S N P		
AAC GAA CCA TCA AAC CCA AAT GAA CCA TCA AAC CCA AAA AAG CCA TCA AAC CCA AAT GAG		900
N E P S N P N E P S N P K K P S N P N E		
CCA TCA AAC CCA AAT GAG CCA TTA AAC CCA AAT GAG CCA TCA AAC CCA AAC GAA CCA TCA		960
P S N P N E P L N P N E P S N P N E P S		
AAC CCA AAC GAA CCA TCA AAC CCA GAA GAA CCA TCA AAC CCT AAA GAG CCA TCA AAC CCA		1020
N P N E P S N P E E P S N P K E P S N P		
AAC GAA CCA TCA AAC CCA GAA GAG CCA AAC CCA GAA GAA CCA TCA AAC CCT AAA GAG CCA		1080
N E P S N P E E P N P E E P S N P K E P		
TCA AAC CCA GAA GAG CCA ATA AAC CCA GAA GAA CTA AAC CCA AAA GAG CCA TCA	-----6-MER REPEAT REGION----->	1140
S N P E E P I N P E E L N P K E P S N P		
GAA GAA TCG AAC CCC AAG GAG CCA ATA AAC CCA GAA GAA TCG AAC CCC AAA GAG CCA ATA		1200
E E S M P K E P I M P E E S M P K E P I		
AAC CCA GAA GAT AAT GAA AAT CCA TTG ATA ATA CAA GAT GAA CCT ATA GAA CCC AGA AAT		1260
N P E D M E M P L I I Q D E P I E P R M		
GAT TCA AAT GTA ATA CCA ATT TTA CCT ATC ATC CCA CAA AAG GGT AAT AAT ATC CCA AGC		1320
D S N V I P I L P I I P Q K G M M I P S		
AAT CTA CCA GAA AAT CCA TCT GAC TCA GAA GTA GAA TAT CCA AGA CCA AAT GAT AAT GGT	<-----(-)-----	1380
M L P E M P S D S E V E Y P R P N D M G		
GAA AAT TCA AAT AAT ACT ATG AAA TCA AAA AAA AAT ATA CCC AAC GAG CGG		1431
E M S M M T M K S K K M I P N E R		




## Results

**Sequence analysis of the 140 kDa antigen gene.** A clone designated  $\lambda$ gMSY-4 (M4) was selected based on its recognition by NYS4. The primary DNA and deduced amino acid sequences of a portion of the genomic DNA insert from clone M4 are shown in Fig. 1(a). Like most other plasmodial antigens, the M4 peptide contains regions of repeating amino acid sequences. One region consists of a repeating trimer Pro-Asn-Asn and the other is composed of, principally, the hexamer Asn-Pro-Asn-Glu-Pro-Ser. The first 48 amino acids are compared in Fig. 1(b) to several proteins that share a remarkable sequence similarity to this region of the M4 peptide. The similarity between region II of the CS protein (301-315) and thrombospondin is well known (25). In addition, mouse properdin was recently shown to contain a similar sequence (26). The M4 peptide shares this conserved motif but contains a more extensive region

of sequence similarity with the recently described TRAP (thrombospondin-related anonymous protein) from *P. falciparum* (27). A search of the PIR protein sequence databank, conducted on 12 February 1990, revealed no additional significant similarity to previously described protein sequences.

**mRNA for the 140 kDa antigen is present in sporozoites.** To demonstrate sporozoite expression of the mRNA for the 140 kDa antigen we utilized the polymerase chain reaction (PCR) specifically primed with M4 clone sequence-based primers (Fig. 1(b)). An amplification product of the predicted size (the distance between the two primers is 1.3 kB) was clearly visible in reaction mixtures which contained, as templates, sporozoite cDNA or DNA from clone M4 (positive control) (Fig. 2, left). In Fig. 2, right, the identity of the amplified cDNA subfragment was confirmed with the M4 probe. A faint signal at 1.3 kB

Fig. 2. Expression of the 140 kDa antigen mRNA in sporozoites as determined by PCR amplification of a cDNA subfragment from reverse-transcribed RNA isolated from infected mosquitoes. Primers used in the amplification are shown in Fig. 1. *Left:* agarose gel stained with ethidium bromide. *Right:* Southern blot of the stained agarose gel probed with  $^{32}$ P-labelled M4 clone DNA. Lanes 1, DNA molecular weight markers; lanes 2, M4 clone DNA (15  $\mu$ g); lanes 3, M4 clone DNA (1.5  $\mu$ g); lanes 4, cDNA (50  $\mu$ g); lanes 5, cDNA (5  $\mu$ g); lanes 6, cDNA (0.5  $\mu$ g); lanes 7, RNA (50  $\mu$ g); lanes 8, RNA (5  $\mu$ g); lanes 9, RNA (0.5  $\mu$ g).



was detected in control reactions of RNA that were not reverse transcribed prior to PCR. We believe this is due to amplification of the genomic sequence which resulted from DNA contamination of the RNA preparation.


**Antibodies to Asn-Pro-Asn-Glu-Pro-Ser recognize the surface of sporozoites and the 140 kDa antigen.** The overall hydrophilicity of the repeating amino acid sequence Asn-Pro-Asn-Glu-Pro-Ser suggested it as a potential antigenic determinant (28). A synthetic peptide containing 3 copies of Asn-Pro-Asn-Glu-Pro-Ser (18-mer) was strongly recognized by NYS4 in an ELISA (data not shown) and mice immunized with the peptide produced antibodies that reacted with sporozoites in an IFAT and with the 140 kDa antigen on Western blots of sporozoite extracts (Fig. 3). These results together with those of Fig. 2 indicate that the M4 peptide sequence corresponds to at least a portion of the 140 kDa antigen and that the antigenic determinant of NYS4 is contained within the repeating hexamer Asn-Pro-Asn-Glu-Pro-Ser. It is notable that this repetitive sequence bears no similarity to the major repeats of the *P. yoelii* CS protein, in which the consensus repeating elements are Gln-Gly-Pro-Gly-Ala-Pro and Gln-Gln-Pro-Pro (29).

## Discussion

In this paper we report the molecular characterization of a portion of the gene encoding what we now call sporozoite surface protein 2 (SSP 2). Like the CS protein, SSP 2 contains an immunogenic sequence of repeating amino acids and the conserved region II domain which is found in a number of other well-characterized proteins. These include thrombospondin, properdin, von Willebrand factor, beta<sub>2</sub>-glycoprotein I, collagen type IV alpha<sub>1</sub> & 2 chains, glycoprotein E from human (alpha) herpesvirus 1, and antistasin, all of them proteins which are thought to play roles in adhesion (30). The sequence similarity between SSP 2 and TRAP, an erythrocytic-stage specific protein of *P. falciparum*, extends beyond the "adhesion motif" of the CS protein in both the amino terminal and carboxyl terminal directions suggesting a more expansive functional domain in these malarial proteins. Nevertheless the conservation of this sequence among proteins from different malarial species underscores its importance to the survival of the parasite. Perhaps sporozoite entry into host cells is facilitated by SSP 2 adherence to extracellular matrices.

Immunization with irradiation-attenuated sporozoites confers sterile immunity against challenge with large numbers of sporozoites. It has long been thought that this protective immunity is directed

Fig. 3. Antibodies to (NPNEPS)<sub>3</sub> recognize the 140 kDa sporozoite surface protein. Western blot of *P. yoelii* sporozoite extract probed with serum from mice immunized with (NPNEPS)<sub>3</sub> conjugated to KLH (lane 1), NYS4 (lane 2), and sera from mice immunized with KLH (lane 3).



entirely against the CS protein. Our findings demonstrate the existence of another protein, SSP 2, on sporozoites that may be an additional target for protective cellular or humoral immune responses.

## References

1. Nussenzweig, V. & Nussenzweig, R.S. Circumsporozoite proteins of malaria parasites. *Cell*, **42**: 401-403 (1985).
2. Cochrane, A.H. et al. Antibody-induced ultrastructural changes of malarial sporozoites. *J. Immunol.*, **116**: 859-867 (1976).
3. Nussenzweig, R.S. et al. Protective immunity produced by the injection of X-irradiated sporozoites of *Plasmodium berghei*. *Nature*, **216**: 160 (1967).

4. Weiss, W.R. et al. CD8<sup>+</sup> cells (cytotoxic/suppressors) are required for protection in mice immunized with malaria sporozoites. *Proc. Natl Acad. Sci., USA*, **85**: 573 (1988).
5. Hoffman, S.L. et al. Immune response gene regulation of immunity to *Plasmodium berghei* sporozoites and circumsporozoite protein vaccines: overcoming genetic restriction with whole organism and subunit vaccines. *J. Immunol.*, **142**: 3581-3584 (1989).
6. Sedegah, M. et al. Use of a vaccinia construct expressing the circumsporozoite protein in the analysis of protective immunity to *Plasmodium yoelii*. In: Lasky, L., ed. *Technological advances in vaccine development*. New York, Alan R. Liss Inc., 1988, pp. 295-309. (UCLA Symposium on Molecular and Cellular Biology, New Series, Vol. 84).
7. Egan, J.E. et al. Efficacy of murine malaria sporozoite vaccines: implications for human vaccine development. *Science*, **236**: 453 (1987).
8. Sadoff, J.C. et al. Oral *Salmonella typhimurium* vaccine expressing circumsporozoite protein protects against malaria. *Science*, **240**: 336-338 (1988).
9. Hoffman, S.L. et al. Sporozoite vaccine induces genetically restricted T-cell elimination of malaria from hepatocytes. *Science*, **244**: 1078-1080 (1989).
10. de la Cruz, V.F. et al. Sequence variation in putative functional domains of the circumsporozoite protein of *Plasmodium falciparum*: implications for vaccine development. *J. biol. chem.*, **262**: 11935-11939 (1987).
11. Good, M.F. et al. Human T-cell recognition of the circumsporozoite protein of *Plasmodium falciparum*: immunodominant T-cell domains map to the polymorphic regions of the molecule. *Proc. Natl Acad. Sci., USA*, **85**: 1199-1203 (1988).
12. Lockyer, M.J. et al. Wild isolates of *Plasmodium falciparum* show extensive polymorphism in T-cell epitopes of the circumsporozoite protein. *Mol. biochem. parasitol.*, **37**: 275-280 (1989).
13. Charoenvit, Y. et al. Characterization of *Plasmodium yoelii* monoclonal antibodies directed against stage-specific sporozoite antigens. *Infect. immun.*, **55**: 604-608 (1987).
14. Wortman, A. et al. Cloning of *Plasmodium yoelii* genes expressing three different sporozoite-specific antigens. *Micro. path.*, **6**: 227-231 (1989).
15. Stephens, R.S. et al. Molecular cloning and expression of *Chlamydia trachomatis* major outer membrane protein antigens in *Escherichia coli*. *Infect. immun.*, **47**: 713-718 (1985).
16. Young, R.A. & Davis, R.W. Efficient isolation of genes by using antibody probes. *Proc. Natl Acad. Sci., USA*, **80**: 1194-1198 (1983).
17. Young, R.A. & Davis, R.W. Yeast RNA polymerase II genes: isolation with antibody probes. *Science*, **222**: 778-782 (1983).
18. Johnson, D.A. et al. Improved technique utilizing non-fat dry milk for analysis of proteins and nucleic acids transferred to nitrocellulose. *Gene anal. techn.*, **1**: 3-4 (1984).
19. Maniatis, T. et al. *Molecular cloning, a laboratory manual*. Cold Spring Harbor, Cold Spring Harbor Laboratory, 1982.
20. Messing, J. New M13 vectors for cloning. *Methods enzymol.*, **101**: 20-78 (1983).
21. Henikoff, S. Unidirectional digestion with exonuclease III in DNA sequence analysis. *Methods enzymol.*, **155**: 156-165 (1987).
22. Chirgwin, J.M. et al. Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry*, **18**: 5294-5299 (1979).
23. Salki, R.K. et al. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science*, **230**: 1350-1354 (1985).
24. Merrifield, B. Peptide synthesis. I. The synthesis of a tetrapeptide. *J. Am. Chem. Soc.*, **85**: 2149-2154 (1963).
25. Lawler, J. & Hynes, R.O. The structure of human thrombospondin, an adhesive glycoprotein with multiple calcium-binding sites and homologies with several different proteins. *J. cell biol.*, **103**: 1635-1648 (1986).
26. Goundis, D. & Reid K.B. Properdin, the terminal complement components, thrombospondin and the circumsporozoite protein of malaria parasites contain similar sequence motifs. *Nature*, **335**: 82-85 (1988).
27. Robson, K.J.H., et al. A highly conserved amino-acid sequence in thrombospondin, properdin and in proteins from sporozoites and blood stages of a human malaria parasite. *Nature*, **335**: 79-82 (1988).
28. Hopp, T.P. & Woods, K.R. Prediction of protein antigenic determinants from amino acid sequences. *Proc. Natl Acad. Sci., USA*, **78**: 3824-3828 (1981).
29. Lal, A.A. et al. Structure of the gene encoding the circumsporozoite protein of *Plasmodium yoelii*. *J. biol. chem.*, **262**: 2937-2940 (1987).
30. Holt, G.D. et al. Antistasin, an inhibitor of coagulation and metastasis, binds to sulfatide (Gal(3-SO<sub>4</sub>) $\beta$ 1-1Cer) and has sequence homology with other proteins that bind sulfated glycoconjugates. *J. biol. chem.*, **264**: 12138-12140 (1989).